

## The Determination of Acid Content in Vinegar

Reading assignment: Chang, Chemistry 10<sup>th</sup> edition, pages 153-156.

### Goals

We will use a titration to determine the concentration of acetic acid in a sample of vinegar in order to become familiar with acid-base reactions.

### Equipment and Supplies

Digital analytical balance, 100-mL beakers (2), 10-mL graduated cylinder, 100-mL volumetric flask, 10-mL graduated pipette, pipette pump, 125-mL Erlenmeyer flasks (3), 50-mL buret, phenolphthalein indicator, ~0.1M sodium hydroxide, vinegar.

Safety Note: Safety glasses are required when performing this experiment.

### Discussion

Acetic acid ( $\text{HC}_2\text{H}_3\text{O}_2$ ) is the active ingredient in vinegar and is responsible for its sour taste. Acetic acid is an example of a weak acid. For a 0.1 mol/L solution of acetic acid only about 1% of the acid ionizes. Compare this to a strong acid like hydrochloric acid. Very close to 100% of hydrochloric acid ionizes. A few examples of strong and weak acids are shown below:


strong acids		weak acids			
perchloric	$\text{HClO}_4$	chlorous	$\text{HClO}_2$	hypochlorous	$\text{HClO}$
sulfuric	$\text{H}_2\text{SO}_4$	hydrocyanic	$\text{HCN}$		
hydrochloric	$\text{HCl}$	sulfurous	$\text{H}_2\text{SO}_3$		
hydrobromic	$\text{HBr}$	hydrosulfuric	$\text{H}_2\text{S}$		
hydroiodic	$\text{HI}$	hydrofluoric	$\text{HF}$		
nitric	$\text{HNO}_3$	nitrous	$\text{HNO}_2$		
		phosphoric	$\text{H}_3\text{PO}_4$		
		acetic	$\text{HC}_2\text{H}_3\text{O}_2$		

Only one of the hydrogen atoms of the acetic acid molecule is acidic:




The structural formula for acetic acid is shown to the right.

The hydrogen attached to the oxygen atom is acidic while the other hydrogen atoms are not.




Another way of representing the acidity of acetic acid is to show its reaction with water:



Here the acetic acid protonates (transfers a proton to) the water molecule. In fact, the hydrogen ion ( $\text{H}^+$ ) is very reactive and doesn't exist in water. However, there is evidence that the hydronium ion does exist. Sometimes equation 1 is used because of its simplicity.

To determine the amount of acetic acid in vinegar (typically 4-5% by mass) we will use an acid-base titration (neutralization reaction). In this experiment we titrate acetic acid with sodium hydroxide (a strong base). The reaction of acetic acid with sodium hydroxide is shown below:




In the reaction between acetic acid and sodium hydroxide, the acetic acid donates a proton to the hydroxide ion and acts as an acid. The hydroxide ion accepts a proton and acts as a base. The stoichiometric relationship between acetic acid and sodium hydroxide is 1:1 (from Equation 2). If the number of moles of NaOH used to titrate a sample of acetic acid are known, then the moles of acetic acid in a sample can easily be found.

Evidence of the reaction between acetic acid and sodium hydroxide isn't visible, so we will add an indicator to determine the equivalence point for the titration. A common acid-base indicator is phenolphthalein. Phenolphthalein is colorless in acidic solution, but changes to pink when the solution becomes basic. At the equivalence point, the moles of acetic acid are equal to the moles of sodium hydroxide, and the solution will turn pink as more sodium hydroxide is added. Phenolphthalein is itself a weak acid. So when enough sodium hydroxide has been added to the acetic acid, such that there is no longer any unreacted acetic acid, the sodium hydroxide will then begin reacting with phenolphthalein. The reaction of phenolphthalein with sodium hydroxide results in a pink solution.

Buret containing sodium hydroxide solution

Erlenmeyer flask with vinegar and indicator



The picture above shows a titration apparatus. The experiment makes use of a buret that contains the sodium hydroxide solution. An Erlenmeyer flask contains the vinegar solution and a couple of drops of the indicator (phenolphthalein). The sodium hydroxide solution is poured into the vinegar solution one drop at a time. The volume of sodium hydroxide required to react with all of the acetic acid in the vinegar is measured from the buret.

### Sample Calculation

A 25.00 mL sample of a hydrofluoric acid, a monoprotic acid, is titrated with a 0.155 M solution of sodium hydroxide. It is found that the indicator changes color once 31.25 mL of the sodium hydroxide solution has been added to the acid. What is the molarity of the acid?

The equation for the reaction is  $\text{HF}(\text{aq}) + \text{NaOH}(\text{aq}) \rightarrow \text{NaF}(\text{aq}) + \text{H}_2\text{O}(\text{l})$

At the equivalence point the moles of the acid (HF) are equal to the moles of the base (NaOH).<sup>\*</sup> We can use the known concentration and measured volume of the sodium hydroxide to find the number of moles of hydroxide used in the titration:


$$\text{moles NaOH} = \left( \frac{0.155 \text{ mol NaOH}}{1 \text{ L}} \right) (0.03125 \text{ L NaOH}) = 0.00484 \text{ mol NaOH}$$
 where the volume has been converted to liters.

At the equivalence point: mol HF = mol NaOH so moles of HF = 0.00484 mol


The concentration of the acid is equal to the number of moles divided by its volume:

$$\text{molarity of acid} = \frac{0.00484 \text{ mol HF}}{0.02500 \text{ L}} = \mathbf{0.194 \text{ mol/L}}$$


<sup>\*</sup>The solution changes color once all of the acid has reacted with the base and some of the indicator reacts with the base. So the change in color often occurs at what is called the endpoint, a little beyond the equivalence point.



125-mL Erlenmeyer flask



50-mL buret



volumetric flask

### SAFETY PRECAUTIONS

Acetic acid and sodium hydroxide are both irritants. Safety glasses must be worn at all times during this experiment. Students perform titrations individually.

1. Obtain about 60 mL of sodium hydroxide (NaOH) in a clean dry beaker. Be sure to write down the concentration of the sodium hydroxide from the bottle (or from the board at the front of the room).
2. Prepare the buret by rinsing the buret with tap water, then with distilled or deionized water, and finally with about 5 mL of sodium hydroxide solution.
3. Fill the buret with sodium hydroxide solution to a level slightly above the 0 mL mark. Open the valve on the buret and allow a couple milliliters to flow into a waste beaker so that any air bubbles can be forced through. Record the volume of the sodium hydroxide in the buret on the data sheet (initial volume).
4. The vinegar we will be using has an acetic acid concentration that would require a large volume of sodium hydroxide for this titration. So we will dilute the vinegar with water before performing the titration. Obtain 15-20 mL of vinegar in a clean and dry 100 mL beaker. Record the brand name of the vinegar.
5. Using a 10-mL volumetric pipette carefully pipet 10 mL of vinegar into a 100-mL volumetric flask.
6. Fill the flask to the 100 mL mark using distilled or deionized water. Use this diluted vinegar to perform the titrations.
7. Measure the mass of 3 clean 125-mL Erlenmeyer flasks (individually) on the digital analytical balance to the nearest 0.1 mg (0.0001 g). Record each mass on the data sheet.
8. Using a 10 mL-graduated cylinder add about 10 mL of the diluted vinegar into each of the weighed Erlenmeyer flasks.
9. Measure the mass of each of the Erlenmeyer flasks containing the diluted vinegar. Record these masses on the data sheet.
10. Add two drops of phenolphthalein indicator to each Erlenmeyer flask containing dilute vinegar and begin the titration by slowly adding the sodium hydroxide solution drop by drop from the buret to the flask. The solution should initially be clear, with no pink tint. Be sure to swirl the solution in the flask

while adding the sodium hydroxide solution. The endpoint for this titration is a very faint pink color that persists for more than 15 seconds. Darker pink colors indicate that you have added too much sodium hydroxide. Read the volume to 0.02 mL.

11. Perform the titration a minimum of three times until you have three sodium hydroxide volumes that are within 0.20 mL of each other. You can perform additional titrations if your delivered volumes are not close to one another.

12. Waste can be disposed of in the sink. Clean your work area before beginning calculations.

Perform these calculations on a separate sheet of paper.

1. Determine the number of moles of sodium hydroxide required to titrate the vinegar for each titration from the known molarity and the titration volume ( $V = V_2 - V_1$ ) of sodium hydroxide. Be sure that the volume of the sodium hydroxide has been converted from milliliters to liters (1 L = 1000 mL).

$$M_{\text{NaOH}} = \frac{\text{moles NaOH}}{\text{Volume NaOH}} \text{ so } \text{moles NaOH} = (V_{\text{NaOH}}) \times (M_{\text{NaOH}})$$

2. The moles of acetic acid are equal to the moles of sodium hydroxide at the equivalence point. The equivalence point is close to the endpoint so we can use the endpoint value. The endpoint is when the phenolphthalein changes color.

$$\text{moles acetic acid} = \text{moles sodium hydroxide}$$

3. Determine the mass of acetic acid present in each titration from the molar mass (sometimes called molecular weight) of acetic acid and moles of acetic acid.

$$\text{molar mass} = \frac{\text{mass}}{\text{moles}} \text{ so } \text{mass acetic acid} = (\text{molar mass}) \times (\text{moles})$$

4. The mass of the vinegar for each titration is found from the measurements using the analytical balance ( $m_2 - m_1$ ).

5. The percentage mass of acetic acid in the vinegar is found from the mass of acetic acid and the mass of vinegar.

$$\% \text{ mass} = \frac{\text{mass of acetic acid}}{\text{mass of vinegar}} \times 100$$

6. The vinegar was diluted by a factor of 10 in this experiment. So the mass percentage of the dilute vinegar should be multiplied by ten to find the mass percentage of the undiluted vinegar.

7. Calculate the average mass percent from your data. You may have one number that seems different from the other two. If so, perform a Q test on your mass percent values to determine if the “suspect” value (outlier) can be rejected. The Q test is performed in the following manner. First calculate Q by finding the ratio of the gap to the range for your data. The gap is the difference between the suspected outlier and the value closest to it. The range is the largest difference between any two numbers from your data. Compare the calculated Q to the table value of Q for the number of data points in your experiment (N). For this experiment, N is usually 3 because three titrations were performed. If the calculated Q value is greater than the Q value listed in the table, then the suspect data point can be rejected and the average is found using the remaining values. If Q calculated is less than Q from the table, then the suspect value must be included in the average.

$$Q_{\text{calc}} = \frac{|\text{gap}|}{|\text{range}|}$$

N	3	4	5
Q*	0.970	0.829	0.710

\*95% confidence

Reject suspect value if  $Q_{\text{calc}} > Q_{\text{table}}$

---

**Observations and Notes**  
**The Determination of Acid Content in Vinegar**

Date \_\_\_\_\_

---

**Data Sheet**  
**Determination of Acid Content in Vinegar**

Name \_\_\_\_\_

Concentration of NaOH \_\_\_\_\_ mol/L

Brand name of vinegar \_\_\_\_\_

	<b>Trial 1</b>	<b>Trial 2</b>	<b>Trial 3</b>
Mass of empty Erlenmeyer flask ( $m_1$ )	_____ g	_____ g	_____ g
Mass of flask and vinegar ( $m_2$ )	_____ g	_____ g	_____ g
Mass of vinegar	_____ g	_____ g	_____ g
Final buret volume ( $V_2$ )	_____ mL	_____ mL	_____ mL
Initial buret volume ( $V_1$ )	_____ mL	_____ mL	_____ mL
Volume of NaOH Delivered ( $V_2 - V_1$ )	_____ mL	_____ mL	_____ mL
Volume of NaOH Delivered in liters	_____ L	_____ L	_____ L
Moles of NaOH used	_____	_____	_____
Moles of acetic acid in sample	_____	_____	_____
Mass of acetic acid	_____ g	_____ g	_____ g
Mass percent (diluted)	_____	_____	_____
Mass percent (undiluted)	_____	_____	_____
Average mass percent (Consider Q test)	_____	_____	_____

---

**Calculations Sheet**  
**Determination of Acid Content in Vinegar**